

□ Appendix 3 □

72 articles which were used in the meta-analysis

- 1) Tsolaki AC, Kosmidou V, Kompatsiaris IY, Papadaniil C, Hadjileontiadis L, Adam A, et al. Brain source localization of MMN and P300 ERPs in mild cognitive impairment and Alzheimer's disease: A high-density EEG approach. *Neurobiol Aging* 2017;55:190-201.
- 2) Tarkka IM, Lehtovirta M, Soininen H, Pääkkönen A, Karhu J, Partanen J. Auditory adaptation is differentially impaired in familial and sporadic Alzheimer's disease. *Biomed Pharmacother* 2002;56(1):45-9.
- 3) Vandoolaeghe E, van Hunsel F, Nuyten D, Maes M. Auditory event related potentials in major depression: Prolonged P300 latency and increased P200 amplitude. *J Affect Disord* 1998;48(2-3):105-13.
- 4) Yamaguchi S, Tsuchiya H, Yamagata S, Toyoda G, Kobayashi S. Event-related brain potentials in response to novel sounds in dementia. *Clin Neurophysiol* 2000;111(2):195-203.
- 5) Sumi N, Nan'no H, Fujimoto O, Ohta Y, Takeda M. Interpeak latency of auditory event-related potentials (P300) in senile depression and dementia of the Alzheimer type. *Psychiatry Clin Neurosci* 2000;54(6):679-84.
- 6) Muscoso EG, Costanzo E, Daniele O, Maugeri D, Natale E, Caravaglios G. Auditory event-related potentials in subcortical vascular cognitive impairment and in Alzheimer's disease. *J Neural Transm (Vienna)* 2006;113(11):1779-86.
- 7) Masanaka T, Hisao T, Keita K, Yasunobu K, Hiroo Y, Bungo O. Multi-modal evoked potentials in corticobasal degeneration, progressive supranuclear palsy and Alzheimer disease. *Int Congr* 2005;1278:145-8.
- 8) Lai CL, Lin RT, Liou LM, Liu CK. The role of event-related potentials in cognitive decline in Alzheimer's disease. *Clin Neurophysiol* 2010; 121(2):194-9.
- 9) Näätänen R, Kujala T, Kreegipuu K, Carlson S, Escera C, Baldeweg T, et al. The mismatch negativity: an index of cognitive decline in neuropsychiatric and neurological diseases and in ageing. *Brain* 2011;134(12):3435-53.
- 10) Hirata K, Hozumi A, Tanaka H, Kubo J, Zeng XH, Yamazaki K, et al. Abnormal information processing in dementia of Alzheimer type. A study using the event-related potential's field. *Eur Arch Psychiatry Clin Neurosci* 2000;250(3):152-5.
- 11) Gungor HA, Yildiz A, Aydin F, Gungor F, Boz A, Ozkaynak S. Tc-99m HMPAO brain SPECT findings in mild and moderate Alzheimer's disease: Correlation with event related potentials. *J Neurol Sci* 2005;234(1-2):47-53.
- 12) Caravaglios G, Costanzo E, Palermo F, Muscoso EG. Decreased amplitude of auditory event-related delta responses in Alzheimer's disease. *Int J Psychophysiol* 2008;70(1):23-32.
- 13) Tachibana H, Takeda M, Okuda B, Kawabata K, Nishimura H, Kodama N, et al. Multimodal evoked potentials in Alzheimer's disease and Binswanger's disease. *J Geriatr Psychiatry Neurol* 1996;9(1):7-12.
- 14) Holt LE, Raine A, Pa G, Schneider LS, Henderson VW, Pollock VE. P300 topography in Alzheimer's disease. *Psychophysiology* 1995;32(3): 257-65.
- 15) Taguchi S, Fujimoto O, Takamori N, Hanatani T, Sumi N, Nan'no H, et al. Neurophysiological evaluation of late paraphrenia: Comparison with chronic schizophrenia and dementia of the Alzheimer type. *Psychogeriatrics* 2003;3(1):29-38.
- 16) van Deursen JA, Vuurman EF, Smits LL, Verhey FR, Riedel WJ. Response speed, contingent negative variation and P300 in Alzheimer's disease and MCI. *Brain Cogn* 2009;69(3):592-9.
- 17) O'Mahony D, Coffey J, Murphy J, O'Hare N, Hamilton D, Rowan M, et al. Event-related potential prolongation in Alzheimer's disease signifies frontal lobe impairment: Evidence from SPECT imaging. *J Gerontol A Biol Sci Med Sci* 1996;51(3):M102-7.
- 18) Bennis K, Portet F, Touchon J, Rondouin G. Diagnostic value of event-related evoked potentials N200 and P300 subcomponents in early diagnosis of Alzheimer's disease and mild cognitive impairment. *J Clin Neurophysiol* 2007;24(5):405-12.
- 19) Fruehwirt W, Dorffner G, Roberts S, Gerstgrasser M, Grossegger D, Schmidt R, et al. Associations of event-related brain potentials and Alzheimer's disease severity: A longitudinal study. *Prog Neuropsychopharmacol Biol Psychiatry* 2019;92:31-38.
- 20) Tanaka F, Kachi T, Yamada T, Sobue G. Auditory and visual event-related potentials and flash visual evoked potentials in Alzheimer's disease: Correlations with Mini-Mental State Examination and Raven's Coloured Progressive Matrices. *J Neurol Sci* 1998;156(1):83-8.
- 21) Jervis B, Belal S, Camilleri K, Cassar T, Bigan C, Linden DE, et al. The independent components of auditory P300 and CNV evoked potentials derived from single-trial recordings. *Physiol Meas* 2007;28(8):745-71.
- 22) Filipović SR, Kostić VS. Utility of auditory P300 in detection of presenile dementia. *J Neurol Sci* 1995;131(2):150-5.
- 23) Uemura JI, Hoshiyama M. Variability of P300 in elderly patients with dementia during a single day. *Int J Rehabil Res* 2007;30(2):167-70.
- 24) Katada E, Sato K, Ojika K, Ueda R. Cognitive event-related potentials: Useful clinical information in Alzheimer's disease. *Current Alzheimer Research* 2004;1(1):63-9.
- 25) Pedroso RV, Cancela JM, Ayán C, Stein AM, Fuzaro G, Costa JLR, et al. Effects of physical exercise on the P300 of elderly with Alzheimer's disease. *J Phys Act Health* 2018;15(6):403-10.
- 26) Wang P, Zhang H, Han L, Zhou Y. Cortical function in Alzheimer's disease and frontotemporal dementia. *Transl Neurosci* 2016;7(1):116-25.
- 27) Knott V, Mohr E, Mahoney C, Engeland C, Ilivitsky V. Effects of acute nicotine administration on cognitive event-related potentials in tacrinetreated and non-treated patients with Alzheimer's disease. *Neuropsychobiology* 2002;45(3):156-60.
- 28) Bokura H, Yamaguchi S, Kobayashi S. Event-related potentials for response inhibition in Parkinson's disease. *Neuropsychologia* 2005;43(6): 967-75.
- 29) Parra MA, Ascencio LL, Urquina HF, Manes F, Ibáñez AM. P300 and neuropsychological assessment in mild cognitive impairment and Alzheimer dementia. *Front Neurol* 2012;3:172.
- 30) Szeliés B, Mielke R, Grond M, Heiss WD. P300 in Alzheimer's disease: Relationships to dementia severity and glucose metabolism. *J Neurol Sci* 1995;130(1):77-81.
- 31) Pokryszko-Dragan A, Słotwiński K, Podemski R. Modality-specific changes in P300 parameters in patients with dementia of the Alzheimer

- type. *Med Sci Monit* 2003;9(4):CR130-4.
- 32) Onofrij M, Thomas A, Luciano AL, Iacono D, Di Rollo A, D'Andreamatteo G, et al. Donepezil versus vitamin E in Alzheimer's disease: Part 2: mild versus moderate-severe Alzheimer's disease. *Clin Neuropharmacol* 2002;25(4):207-15.
 - 33) Mochizuki Y, Oishi M, Takasu T. Correlations between P300 components and regional cerebral blood flows. *J Clin Neurosci* 2001;8(5):407-10.
 - 34) Lee MS, Lee SH, Moon EO, Moon YJ, Kim S, Kim SH, et al. Neuropsychological correlates of the P300 in patients with Alzheimer's disease. *Prog Neuropsychopharmacol Biol Psychiatry* 2013;40:62-9.
 - 35) Juckel G, Clotz F, Frodl T, Kawohl W, Hampel H, Pogarell O, et al. Diagnostic usefulness of cognitive auditory event-related p300 subcomponents in patients with Alzheimers disease? *J Clin Neurophysiol* 2008;25(3):147-52.
 - 36) Yamamoto M, Kachi T, Igata A. Pain-related somatosensory evoked potentials in dementia. *J Neurol Sci* 1996;137(2):117-9.
 - 37) Jiménez-Escrig A, Fernández-Lorente J, Herrero A, Baron M, Lousa M, de Blas G, et al. Event-related evoked potential P300 in frontotemporal dementia. *Dement Geriatr Cogn Disord* 2002;13(1):27-32.
 - 38) Gironell A, García-Sánchez C, Estévez-González A, Boltes A, Kulisevsky J. Usefulness of p300 in subjective memory complaints: A prospective study. *J Clin Neurophysiol* 2005;22(4):279-84.
 - 39) Bonanni L, Franciotti R, Onofrij V, Anzellotti F, Mancino E, Monaco D, et al. Revisiting P300 cognitive studies for dementia diagnosis: Early dementia with Lewy bodies (DLB) and Alzheimer disease (AD). *Neurophysiol Clin* 2010;40(5-6):255-65.
 - 40) Claus JJ, Mohr E. Attentional deficits in Alzheimer's, Parkinson's, and Huntington's diseases. *Acta Neurol Scand* 1996;93(5):346-51.
 - 41) Frodl T, Hampel H, Juckel G, Bürger K, Padberg F, Engel RR, et al. Value of event-related P300 subcomponents in the clinical diagnosis of mild cognitive impairment and Alzheimer's disease. *Psychophysiology* 2002;39(2):175-81.
 - 42) Boller F, El Massioui F, Devouche E, Traykov L, Pomati S, Starkstein SE. Processing emotional information in Alzheimer's disease: Effects on memory performance and neurophysiological correlates. *Dement Geriatr Cogn Disord* 2002;14(2):104-12.
 - 43) Ashford JW, Coburn KL, Rose TL, Bayley PJ. P300 energy loss in aging and Alzheimer's disease. *J Alzheimers Dis* 2011;26 Suppl 3:229-38.
 - 44) Ally BA, Jones GE, Cole JA, Budson AE. The P300 component in patients with Alzheimer's disease and their biological children. *Biol Psychol* 2006;72(2):180-7.
 - 45) Knott V, Mohr E, Haché N, Mahoney C, Mendis T. EEG and the passive P300 in dementia of the Alzheimer type. *Clin Electroencephalogr* 1999;30(2):64-72.
 - 46) Katada E, Sato K, Sawaki A, Dohi Y, Ueda R, Ojika K. Long-term effects of donepezil on P300 auditory event-related potentials in patients with Alzheimer's disease. *J Geriatr Psychiatry Neurol* 2003;16(1):39-43.
 - 47) Bender S, Bluschke A, Dippel G, Rupp A, Weisbrod M, Thomas C. Auditory post-processing in a passive listening task is deficient in Alzheimer's disease. *Clin Neurophysiol* 2014;125(1):53-62.
 - 48) Missonnier P, Ragot R, Derouesné C, Guez D, Renault B. Automatic attentional shifts induced by a noradrenergic drug in Alzheimer's disease: Evidence from evoked potentials. *Int J Psychophysiol* 1999;33(3):243-51.
 - 49) Croot K, Hodges JR, Patterson K. Evidence for impaired sentence comprehension in early Alzheimer's disease. *J Int Neuropsychol Soc* 1999;5(5):393-404.
 - 50) Golden HL, Agustus JL, Goll JC, Downey LE, Mummery CJ, Schott JM, et al. Functional neuroanatomy of auditory scene analysis in Alzheimer's disease. *Neuroimage Clin* 2015;7:699-708.
 - 51) Golden HL, Agustus JL, Nicholas JM, Schott JM, Crutch SJ, Mancini L, et al. Functional neuroanatomy of spatial sound processing in Alzheimer's disease. *Neurobiol Aging* 2016;39:154-64.
 - 52) White DA, Murphy CF. Working memory for nonverbal auditory information in dementia of the Alzheimer type. *Arch Clin Neuropsychol* 1998;13(4):339-47.
 - 53) Rösler A, Mapstone M, Hays-Wicklund A, Gitelman DR, Weintraub S. The "zoom lens" of focal attention in visual search: Changes in aging and Alzheimer's disease. *Cortex* 2005;41(4):512-9.
 - 54) Perry RJ, Hodges JR. Fate of patients with questionable (very mild) Alzheimer's disease: Longitudinal profiles of individual subjects' decline. *Dement Geriatr Cogn Disord* 2000;11(6):342-9.
 - 55) Manuel AL, Foxe D, Bradshaw N, Cordato NJ, Hodges JR, Burrell JR, et al. Sustained attention failures on a 3-min reaction time task is a sensitive marker of dementia. *J Neurol* 2019;266(6):1323-31.
 - 56) Hardy CJD, Marshall CR, Bond RL, Russell LL, Dick K, Ariti C, et al. Retained capacity for perceptual learning of degraded speech in primary progressive aphasia and Alzheimer's disease. *Alzheimers Res Ther* 2018;10(1):70.
 - 57) Sommers MS. Spoken word recognition in individuals with dementia of the Alzheimer's type: Changes in talker normalization and lexical discrimination. *Psychol Aging* 1998;13(4):631-46.
 - 58) Waters G, Caplan D. Working memory and online syntactic processing in Alzheimer's disease: Studies with auditory moving window presentation. *J Gerontol B Psychol Sci Soc Sci* 2002;57(4):P298-311.
 - 59) Strouse AL, Hall JW 3rd. Test-retest reliability of a dichotic digits test for assessing central auditory function in Alzheimer's disease. *Audiology* 1995;34(2):85-90.
 - 60) Gates GA, Anderson ML, Feeney MP, McCurry SM, Larson EB. Central auditory dysfunction in older persons with memory impairment or Alzheimer dementia. *Archives of Otolaryngology-Head & Neck Surgery* 2008;134(7):771-7.
 - 61) Gates GA, Karzon RK, Garcia P, Peterein J, Storandt M, Morris JC, et al. Auditory dysfunction in aging and senile dementia of the Alzheimer's type. *Arch Neurol* 1995;52(6):626-34.
 - 62) Strouse AL, Hall JW 3rd, Burger MC. Central auditory processing in Alzheimer's disease. *Ear Hear* 1995;16(2):230-8.
 - 63) Saxton J, Munro CA, Butters MA, Schramke C, McNeil MA. Alcohol, dementia, and Alzheimer's disease: Comparison of neuropsychological profiles. *J Geriatr Psychiatry Neurol* 2000;13(3):141-9.
 - 64) Pachana NA, Boone KB, Miller BL, Cummings JL, Berman N. Comparison of neuropsychological functioning in Alzheimer's disease and frontotemporal dementia. *J Int Neuropsychol Soc* 1996;2(6):505-10.
 - 65) Loring DW, Goldstein FC, Chen C, Drane DL, Lah JJ, Zhao L, et al. False-positive error rates for reliable digit span and auditory verbal learning test performance validity measures in amnesic mild cognitive impairment and early Alzheimer disease. *Arch Clin Neuropsychol*

- 2016;31(4):313-31.
- 66) Luzzi S, Baldinelli S, Ranaldi V, Fabi K, Cafazzo V, Fringuelli F, et al. Famous faces and voices: Differential profiles in early right and left semantic dementia and in Alzheimer's disease. *Neuropsychologia* 2017;94:118-28.
 - 67) Thompson SB, Macdonald J, Coates TD. Improving visual memory with Aricept (donepezil hydrochloride, E2020) in mild-to-moderate Alzheimer's disease. *Clin Gerontol* 2001;24(1-2):55-73.
 - 68) Ruzzoli M, Pirulli C, Mazza V, Miniussi C, Brignani D. The mismatch negativity as an index of cognitive decline for the early detection of Alzheimer's disease. *Sci Rep* 2016;6:33167.
 - 69) Xiao S, Yao P, Li X, Zhang M. Neuropsychological testing profiles of patients with Alzheimer's disease and mild cognitive impairment: A case-control study. *Hong Kong J Psychiatry* 2002;12(4):2-5.
 - 70) Thomas DA, Libon DJ, Ledakis GE. Treating dementia patients with vascular lesions with donepezil: A preliminary analysis. *Appl Neuropsychol* 2005;12(1):12-8.
 - 71) Libon DJ, Mattson RE, Glosser G, Kaplan E, Malamut BL, Sands LP, et al. A nine—word dementia version of the California verbal learning test. *Clin Neuropsychol* 1996;10(3):237-44.
 - 72) Zhao Q, Lv Y, Zhou Y, Hong Z, Guo Q. Short-term delayed recall of auditory verbal learning test is equivalent to long-term delayed recall for identifying amnesic mild cognitive impairment. *PLoS One* 2012;7(12):e51157.

정답 및 해설

답 ④

해설 증례는 전두동에 발생한 골종으로 부비동에서 가장 흔히 발견되는 양성종양이다.

참고 문헌: 대한이비인후과학회. 이비인후과학:비과. 개정2판. 파주: 군자출판사;2018. p.544-5.